


TRULICITY® - SUMMARY OF IMPORTANT INFORMATION


Where to inject ?

- Your doctor, pharmacist or nurse can help you choose the injection site that is best for you.
- You may inject the medicine into your stomach (abdomen) or thigh.
- Another person may give you the injection in your upper arm.
- Change (rotate) your injection site each week. You may use the same area of your body, but be sure to choose a different injection site in that area.



Where to dispose of the pen?

- Dispose of the Trulicity® pen with your normal household waste.



What is GLP-1 and how does Trulicity® work?


GLP-1 is a natural hormone that is released when food is eaten. If you have type 2 diabetes, your body may not produce sufficient amounts of this hormone.

- It tells your **pancreas** how much insulin to release after a meal.
- It can limit how much sugar gets from your **liver** into your blood.
- It reduces the speed at which food leaves your **stomach**.
- It can reduce your appetite (although Trulicity® is not a weight-loss product).
- These effects can help lower your blood sugar levels. A GLP-1 receptor agonist mimics this effect.

Where to store?

- Store your Pen in the refrigerator.
- When refrigeration is not possible, you can keep your Pen at room temperature (below 30°C) for up to a total of 14 days.
- For complete information about proper storage, read the Patient Information Leaflet.

HOW TO USE TRULICITY®, STEP BY STEP.




1. UNCAP



grey cap



2. PLACE AND UNLOCK



3. PRESS AND HOLD



- **Make sure the Pen is locked.**
- Remove the grey cap.
- Do not replace the cap. This could damage the needle. Do not touch the needle.
- After you remove it, the grey cap should look like the picture.

If the grey cap does not look like the one pictured above:

- Do **NOT** use the pen.
- Keep the pen and cap safe and contact Lilly.
- Use a new pen.

- Place the clear base of the pen flat and firmly against your skin at the injection site.
- **Unlock** the pen by turning the locking ring.

- Press and hold the green injection button. You will hear a loud click.
- **Continue holding the clear base of the pen firmly against your skin until you hear a second click. This occurs when the needle starts retracting in about 5-10 seconds.**
- Remove the pen from the injection site.

- You will know your injection is complete when the grey part is visible.

OUR SERVICE

Videos on the simple use and mode of action (in different languages) can be found here (password: Zucker):

Watch the videos here



Or call us at this number:
 Telephone: 06172 - 273 2222 Mon. - Fri. 09:00 AM - 3:00PM
 @ med_info@lilly.com

Leben so normal wie möglich | www.lilly-pharma.de
www.lilly-diabetes.de

PLEASE SEE THE INSTRUCTIONS FOR USE FOR FURTHER INFORMATION.

